



# **Travis County Influenza Surveillance**

This report contains data for the 2014-2015 Influenza Season through November 22, 2014 (MMWR<sup>1</sup> week 47).

#### **Situation Update:**

- CDC reports seasonal flu activity increased slightly in the United States and flu activity in Texas rose from minimal to moderate.
- Nationally, the percentage of specimens testing positive for influenza viruses in the United States increased to 12.6%.
- Influenza A (H3N2) and influenza B viruses have all been identified as most common in the U.S. this season. Very few 2009 influenza A (H1N1) positive specimens have been reported.
- Nationally, 1123 (91.4%) of the 1228 influenza-positive tests reported to CDC were influenza A viruses and 105 (8.6%) were influenza B viruses. Of the 354 influenza A viruses that were subtyped, 99.7% were H3 viruses.
- Statewide, 24.85% specimens tested in influenza surveillance labs were positive for influenza; 625 (86.45%) of the 723 influenza-positive tests were influenza A viruses and 98 (13.55%) were influenza B viruses. Of the 115 influenza A viruses that were subtyped, 100% were H3 viruses. Thirteen specimens were submitted in Travis County for influenza testing, twelve of the specimens tested positive for H3N2.
- Nationwide, 2.0% of patient visits reported through the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) were due to influenza-like illness (ILI). This percentage is at the national baseline of 2.0%.
- In Texas, the percentage of visits for influenza-like illness as reported by ILINet providers was 6.62%, a moderate level of ILI activity. Locally, the percentage of visits was 3.87%.
- Four pediatric influenza related deaths have been reported nationally; no pediatric influenza deaths have been reported in Texas this season, none in Travis County.
- Nationwide, the proportion of deaths attributed to pneumonia and influenza is below the epidemic threshold of 6.4%.
   Locally, the number of deaths attributed to pneumonia and influenza has trended near seasonal levels locally. None of these deaths have been attributed to influenza.





#### H3N2v:

- One infection with an influenza A (H3N2) variant virus (H3N2v) was reported by Wisconsin. The patient is reported to
  have had close contact with swine in the week prior to illness onset. No ongoing human-to-human transmission has
  been identified. This is the first H3N2v infection reported for the 2014-2015 influenza season.
- All cases of H3N2v have mostly been associated with prolonged exposure to pigs at agricultural fairs. Limited humanto-human spread of this virus has been detected in the past but no sustained or community spread has been identified.
   CDC reports that it is possible for sporadic infections and even localized outbreaks to continue to occur.
- For more information, visit <u>H3N2v Influenza</u>.

#### **Preventive Actions:**

- The CDC recommends that anyone six months of age and older should be vaccinated for the flu every flu season. Individuals who are at high risk are especially advised to be vaccinated. High risk individuals are as follows:
  - o Persons 65 years of age and older
  - o Pregnant women
  - o Individuals with certain chronic medical conditions, including diabetes, asthma, heart and lung disease
  - Health care workers
  - o Individuals who live with and/or care for high risk individuals
- In addition to the vaccine, there are certain preventive measures an individual can take to prevent the flu:
  - o Wash your hands often with soap and water, or use an alcohol-based hand sanitizer if soap is not available.
  - o Practice proper sneezing and coughing measures.
  - o Avoid touching your nose, mouth and eyes.
  - o Avoid coming into contact with an individual who is sick.
  - o If you are sick, stay home until you are fever free for 24 hours without the use of fever-reducing medications.

Although these preventive actions are highly recommended they do not take the place of the flu vaccine.

Other Helpful Resources: <u>Texas Department of State Health Services (DSHS)</u>

World Health Organization (WHO)

Centers for Disease Control and Prevention (CDC)





Figure 1. Influenza Surveillance, Travis County Providers <sup>4,5,6</sup>

Week Ending	10/4/2014	10/11/2014	10/18/2014	10/25/2014	11/1/2014	11/8/2014	11/15/2014	11/22/2014
MMWR Week	40	41	42	43	44	45	46	47
# Influenza Rapid Tests Performed	139	181	408	372	236	191	575	693
# Total Positive Influenza Tests	7	7	18	18	5	12	47	148
% Positive Influenza Tests	5.0%	3.9%	4.4%	4.8%	2.1%	6.3%	8.2%	21.4%
# Positive A Influenza	6	5	11	14	2	5	35	115
# Positive B Influenza	1	2	7	4	3	7	11	31
# Non-Differentiated Influenza <sup>3</sup>	0	0	0	0	0	0	1	2

Data source: Austin/Travis County Influenza surveillance reporters

Figure 2. Travis County Confirmed Influenza Test Results: 2,9,10

		12/1/2013 - 2/1/20143	2/2/2014 - 4/5/2014	4/6/2014 - 6/7/2014	6/8/2014 - 8/9/2014	8/10/2014 - 10/11/2014	10/12/2014 - 02/14/2015
N	MMWR Week	49-5	6-14	15-23	24-32	25-41	42-6*
(Total F	Influenza A Positive PCR Tests)	91	5	1	0	1	0
Subtype	Seasonal H1N1	83	2	0	0	0	0
	Season H3N2	2	3	1	0	1	0
	Not Subtyped	13	0	0	0	0	0
(Total F	Influenza B Positive PCR Tests)	1	6	0	0	0	0
PCR N	egative Specimens	123	17	4	7	4	6

<sup>\*</sup> Incomplete data.

Data source: Austin/Travis County Influenza surveillance reporters and the Department of State Health Services lab

<sup>\*\* 1</sup> test was culture confirmed rather than PCR confirmed





Figure 3. Influenza Testing by Texas Laboratories 11,14

	Week 47					
Number of labs reporting flu tests	15					
Number of specimens tested	2910					
Number of positive specimens (%) <sup>†</sup>	723 (24.85%)					
Percentage of total tests that were antigen detection tests	68.32%					
Positive specimens by type/subtype [n (%)]						
Influenza A	625 (86.45%)					
Subtyping performed	115 (18.40%)					
A (H1N1)	0 (0.00%)					
A (H3N2)	115 (100.00%)					
Subtyping not performed	510 (81.60%)					
Influenza B	98 (13.55%)					

# Texas Antigenic Characterization<sup>11</sup>

### Influenza A (H3N2) [5]

• Five (100%) viruses were related to A/Texas/50/2012, the influenza A (H3N2) component of the 2014-2015 Northern Hemisphere influenza vaccine.

### Influenza B [3]

- Yamagata lineage [1]: One (100%) influenza B/Yamagata-lineage virus has been characterized from Texas.
   A B/ Massachusetts/2/2012-like virus is included as an influenza B component of the 2014-2015 Northern Hemisphere trivalent and quadrivalent influenza vaccines.
- Victoria lineage [2]: Two (67%) influenza B/Victoria-lineage viruses were characterized as B/Brisbane/60/2008-like, which is included as an influenza B component of the 2014-2015 Northern Hemisphere quadrivalent influenza vaccine.

### Texas Antiviral Resistance<sup>11</sup>

No antiviral resistance testing data for Texas specimens are available at this time.






Figure 4. Texas Respiratory Laboratory Results:

Virus	Number of Laboratories Testing	Tests Performed	Positive Tests	Percentage of Tests Positive
Adenovirus (respiratory)	6	611	18	2.95%
HMPV	5	614	9	1.47%
Parainfluenza virus	7	706	42	5.95%
Rhinovirus	4	378	91	24.07%
RSV <sup>†</sup>	11	1341	242	18.05%

Figure 5. Number Tested and Percent Positive Rapid Influenza Tests by Week, Travis County: 2012-2013 Influenza Season<sup>4,5,6,8</sup>




Data source: Austin/Travis County Influenza surveillance reporters





Figure 4. Percentage of Visits Due to Influenza-like-Illness Reported by Travis County Participants in ILINet: 2009-2013<sup>8</sup>




Data source: CDC ILI.net system





Figure 5. Statewide Influenza Activity<sup>10,12,13</sup>




This chart was obtained from the Texas Influenza Surveillance Report http://www.dshs.state.tx.us/idcu/disease/Influenza/surveillance/2013/






Figure 6. Annual Influenza Related Mortality, Travis County: 1990-2009



Data source: Texas Department of State Health Services http://soupfin.tdh.state.tx.us/deathdoc.htm Data as of September 12, 2012

Figure 7. City of Austin Pneumonia and Influenza Mortality: 2010-2013




Data source: Center for Disease Control and Prevention 122 Cities Mortality: http://wonder.cdc.gov/mmwr/mmwrmort.asp






Figure 10. RODS Syndromic Surveillance for the Respiratory Prodrome Category, **Travis County: 2010-2013** 



Data source: University of Pittsburgh Real-time Outbreak and Disease Surveillance System. Data is an aggregation of Austin area chief complaint hospital data

Figure 11. RODS Syndromic Surveillance for Influenza-Like-Illness, **Travis County: 2010-2013** 







For additional information about Influenza surveillance, contact:
The Austin/Travis County Health and Human Services Epidemiology and Health Statistics Unit at (512) 972-5555

- MMWR is the Morbidity and Mortality weekly report week published by the CDC.
- <sup>2</sup> PCR testing is performed for specimens referred by area sentinel Influenza surveillance reporters. Providers interested in becoming sentinel providers may call the Austin/Travis County Health and Human Services Epidemiology and Health Statistics Unit at (512) 972-5555.
- Influenza A is commonly split into 2 subtypes: H1N1 and H3N2. Both strains can circulate each Influenza season.
- Influenza is not a reportable condition in Texas; therefore, data is provided by sentinel surveillance reporters and is only a sample of the Influenza activity occurring in the Austin/Travis County area.
- Data represent rapid Influenza testing; these tests provide quick results reporting only Influenza A or B (no subtyping). Early in the flu season, results should be used with caution due to false positive results which can occur, especially during times when Influenza activity is low.
- <sup>6</sup> Influenza data is collected from a variety of reporters; the number of reporters can vary from week to week.
- Non-Differentiated refers to rapid test results that do not differentiate between Influenza A and B.
- <sup>8</sup> Data for Austin/Travis County ILI reporters only; the number of reporters can vary from week to week.
- Influenza is confirmed via PCR testing and a further subtyping may be performed. PCR testing is performed for specimens referred by area Influenza surveillance reporters.
- Positive laboratory results are reported according to specimen collection date or date received in the lab if the former is unknown.
- Antigenic characterization and antiviral resistance is obtained from the Texas Influenza Surveillance Report <a href="http://www.dshs.state.tx.us/idcu/disease/Influenza/surveillance/2013/">http://www.dshs.state.tx.us/idcu/disease/Influenza/surveillance/2013/</a>.
- <sup>12</sup> Influenza activity level corresponds to the current MMWR week only and does not reflect the previous weeks' activity.
- 13 The majority of Influenza cases are not reportable by law to the Texas Department of State Health Services. This map contains data from sentinel sites and does not represent all Influenza cases in the state.
- Laboratory data in 2013-2014 season reports may not be comparable to reports from previous seasons because of the inclusion of DSHS and LRN laboratory data for the current season.
- <sup>15</sup> Some non-NREVSS reporters also contribute to the RSV data.