



Austin/Travis County Health and Human Services Department
Epidemiology and Health Statistics Unit



Travis County Influenza Surveillance

This report contains data for the 2013-2014 Influenza Season through March 1, 2014 (MMWR¹ week 9).

Situation Update:

- National flu activity continues to decrease, but remains elevated in certain indicators.
- Influenza like illness (ILI) remains above baseline nationwide; however, it has decreased over the past nine weeks reaching the national baseline of 2.0%. In Texas, the percentage of visits for influenza-like illness as reported by ILINet providers was 5.13% which is a moderate level of ILI activity. Locally, the percentage of visits was 3.74%.
- Nationally, pneumonia and influenza deaths remain above the epidemic threshold (7.9% of total deaths). Locally these deaths have trended near or below seasonal levels and 20 deaths have been attributed to influenza.
- Nationally, there have been 65 pediatric influenza-associated influenza deaths, 11 reported in Texas. There have been two pediatric influenza-associated deaths in Travis County.
- The percentage of specimens testing positive for influenza decreased nationally and is at 8.7%. There was a slight increase statewide; the percent positive is for the week is 13.54%. Locally 0 out of 2 specimens submitted for PCR testing were positive for influenza.
- Of the 587 positive specimens tested nationally, 80% were influenza A and 20% were influenza B. Subtyping was performed on 253 influenza A specimens, 19.4% were H3 and 80.6% were 2009 H1N1. Statewide, influenza A comprises 27.3% of the specimens submitted. Further testing performed on 16 of these specimens show H1N1 accounts for 62.5% of the influenza A.
- Of the 1541 influenza specimens antigenically characterized by the CDC so for this year, 1321 specimens matched the influenza A H1N1 component and 170 specimens matched the influenza A H3N2 component in the trivalent and quadravalent influenza vaccines. Of the 50 influenza B specimens, 31 specimens matched the component found in both trivalent and quadravalent influenza vaccines and the other 19 specimens matched the second B strain of the quadravalent influenza vaccine.
- In Texas, characterization has been performed on 69 influenza A (H1N1) viruses and ten influenza A (H3N2) viruses both matching components of the trivalent and quadravalent influenza vaccines. Additionally, three influenza B viruses have been characterized as one of the B components of the quadravalent vaccine but not in the trivalent vaccine.



**Austin/Travis County Health and Human Services Department
Epidemiology and Health Statistics Unit**



- Some antiviral resistance to oseltamivir (Tamiflu) has been seen in 34 out of 3733 specimens tested (<1%), all of which were 2009 H1N1. High levels of resistance to adamantanes continues to occur among 2009 H1N1 and H3N2 viruses. Adamantanes are not effective against influenza B. Texas has had 330 specimens tested two have been resistant to oseltamivir (Tamiflu).

H5N1:

- Canada reported the first case of influenza A H5N1, in a human, ever detected in the Americas. No cases of H5N1 have been reported in the United States. The CDC reports the risk to people in the United States is considered to be low at this time.
- H5N1 infections are rare and these viruses do not spread easily from person to person. The CDC is not recommending that the public take and special actions regarding H5N1 in response to this case.
- For more information, visit [H5N1 Influenza](#).

H7N9:

- No cases of H7N9 have been reported in the United States. The CDC reports the risk to people in the United States is considered to be low at this time.
- For more information, visit [H7N9 Influenza](#).



Austin/Travis County Health and Human Services Department Epidemiology and Health Statistics Unit



Preventive Actions:

- The CDC recommends that anyone six months of age and older should be vaccinated for the flu every flu season. Individuals who are at high risk are especially advised to be vaccinated. High risk individuals are as follows:
 - Persons 65 years of age and older
 - Pregnant women
 - Individuals with certain chronic medical conditions, including diabetes, asthma, heart and lung disease
 - Health care workers
 - Individuals who live with and/or care for high risk individuals
- In addition to the vaccine, there are certain preventive measures an individual can take to prevent the flu:
 - Wash your hands often with soap and water, or use an alcohol-based hand sanitizer if soap is not available.
 - Practice proper sneezing and coughing measures.
 - Avoid touching your nose, mouth and eyes.
 - Avoid coming into contact with an individual who is sick.
 - If you are sick, stay home until you are fever free for 24 hours without the use of fever-reducing medications.

Although these preventive actions are highly recommended they do not take the place of the flu vaccine.

For additional information about Influenza surveillance, contact:

The Austin/Travis County Health and Human Services Epidemiology and Health Statistics Unit at (512) 972-5555

Other Helpful Resources:

[Texas Department of State Health Services \(DSHS\)](#)

[Centers for Disease Control and Prevention \(CDC\)](#)

[World Health Organization \(WHO\)](#)



**Austin/Travis County Health and Human Services Department
Epidemiology and Health Statistics Unit**




Figure 1. Influenza Surveillance, Travis County Providers ^{4,5,6}

Week Ending	1/11/2014	1/18/2014	1/25/2014	2/1/2014	2/8/2014	2/15/2014	2/22/2014	3/1/2014
MMWR Week	2	3	4	5	6	7	8	9
# Influenza Rapid Tests Performed	2055	1650	1494	1127	962	868	879	684
# Total Positive Influenza Tests	483	401	256	168	156	169	147	105
% Positive Influenza Tests	23.5%	24.3%	17.1%	14.9%	16.2%	19.5%	16.7%	15.4%
# Positive A Influenza	476	387	228	137	99	110	70	47
# Positive B Influenza	6	13	21	26	54	57	72	58
# Non-Differentiated Influenza ³	1	1	7	5	3	2	5	0

Data source: Austin/Travis County Influenza surveillance reporters

Figure 2. Number Tested and Percent Positive Rapid Influenza Tests by Week, Travis County: 2012-2013 Influenza Season ^{4,5,6,8}



Data source: Austin/Travis County Influenza surveillance reporters


Austin/Travis County influenza surveillance does not attempt to capture all cases of influenza or influenza-like illness. These data should be used for trending purposes over time and for identifying types/strains of influenza that are occurring in the Austin area rather than for estimating the total number of cases.



**Austin/Travis County Health and Human Services Department
Epidemiology and Health Statistics Unit**



Figure 3. Percentage of Visits Due to Influenza-like-Illness Reported by Travis County Participants in ILINet: 2009-2013⁸



Data source: CDC ILI.net system

Figure 4. Travis County Confirmed Influenza Test Results:^{2,9,10}

		3/24/2013 - 5/25/2013	05/26/2013 - 7/27/2013	7/28/2013 - 9/28/2013	09/29/2013 - 11/30/2013	12/1/2013 - 2/1/2014	2/2/2014 - 4/5/2014
MMWR Week		13-21	22-30	31-39	40-48	49-5	6-14*
Influenza A (Total Positive PCR Tests)		1	0	1	6	91	4
Subtype	Seasonal H1N1	1	0	1	6	83	2
	Season H3N2	0	0	0	0	2	2
	Not Subtyped	0	0	0	0	13	0
Influenza B (Total Positive PCR Tests)		0	0	0	0	1	1
PCR Negative Specimens		6	0	9	13	123	5

* Incomplete data.

Data source: Austin/Travis County Influenza surveillance reporters and the Department of State Health Services lab

Austin/Travis County influenza surveillance does not attempt to capture all cases of influenza or influenza-like illness. These data should be used for trending purposes over time and for identifying types/strains of influenza that are occurring in the Austin area rather than for estimating the total number of cases.



**Austin/Travis County Health and Human Services Department
Epidemiology and Health Statistics Unit**



Figure 5. Influenza Testing by Texas Laboratories^{11, 14}

Week 09	
Number of labs reporting flu tests	13
Number of specimens tested	1433
Number of positive specimens (%) [†]	194(13.54%)
Percentage of total tests that were antigen detection tests	68.32%
Positive specimens by type/subtype [n (%)]	
Influenza A	53 (27.3%)
Subtyping performed	16 (30.2%)
A (H1N1)	10 (62.5%)
A (H3N2)	6 (37.5%)
Subtyping not performed	37 (69.8%)
Influenza B	141 (72.7%)

[†]Laboratory data in 2013-2014 season reports may not be comparable to reports from previous seasons because of the inclusion of DSHS and LRN laboratory data for the current season.

Figure 6. Texas Antiviral Resistance¹¹

	Oseltamivir		Zanamivir	
	Virus samples tested (n)	Resistant viruses, number (%)	Virus samples tested (n)	Resistant viruses, number (%)
Influenza A (H1N1)	327	2 (0.61%)	13	0 (0%)
Influenza A (H3N2)	3	0 (0%)	3	0 (0%)
Influenza B	0	--	0	--

- Since October 1, 2013, 2 influenza viruses from Texas have shown resistance to Oseltamivir (Tamiflu).

Texas Antigenic Characterization¹¹

Austin/Travis County influenza surveillance does not attempt to capture all cases of influenza or influenza-like illness. These data should be used for trending purposes over time and for identifying types/strains of influenza that are occurring in the Austin area rather than for estimating the total number of cases.



**Austin/Travis County Health and Human Services Department
Epidemiology and Health Statistics Unit**



Influenza A (H1N1) [69]

- Fifty-two viruses were related to A/California/07/2009. This virus strain was included in the 2013-2014 influenza vaccine for the Northern Hemisphere.

Influenza A (H3N2) [10]

- Three viruses were related to A/Texas/50/2012, an A (H3N2) virus antigenically like the cell-propagated prototype virus A/Victoria/361/2011. This virus strain was included in the 2013-2014 influenza vaccine for the Northern Hemisphere.

Influenza B [3]

- Yamagata lineage [0]: No (0%) influenza B/Yamagata-lineage viruses have been characterized from Texas. B/Massachusetts/2/2012-like is included as an influenza B component of the 2013-2014 Northern Hemisphere trivalent and quadrivalent influenza vaccines.
- Victoria lineage [3]: Three (100%) influenza B/Victoria-lineage viruses were characterized as B/Brisbane/60/2008-like, which is included as an influenza B component of the 2013-2014 Northern Hemisphere quadrivalent influenza vaccine.

Figure 7. Texas Respiratory Laboratory Results⁵

Virus	Number of Laboratories Testing	Tests Performed	Positive Tests	Percentage of Tests Positive
Adenovirus (respiratory)	4	346	15	4.34%
HMPV	6	380	35	9.21%
Parainfluenza virus	5	466	5	1.07%
Rhinovirus	3	334	95	28.44%
RSV [†]	11	561	137	24.42%


[†]RSV tests displayed in the table are antigen detection tests only. Some non-NREVSS reporters also contribute to the RSV data.



Austin/Travis County Health and Human Services Department
Epidemiology and Health Statistics Unit



Figure 8. Statewide Influenza Activity^{10,12,13}



This map was obtained from the Texas Influenza Surveillance Report <http://www.dshs.state.tx.us/idcu/disease/Influenza/surveillance/2013/>


Austin/Travis County influenza surveillance does not attempt to capture all cases of influenza or influenza-like illness. These data should be used for trending purposes over time and for identifying types/strains of influenza that are occurring in the Austin area rather than for estimating the total number of cases.



**Austin/Travis County Health and Human Services Department
Epidemiology and Health Statistics Unit**




**Figure 9. Annual Influenza Related Mortality,
Travis County: 1990-2010**



Data source: Texas Department of State Health Services <http://soupin.tdh.state.tx.us/deathdoc.htm> Data as of September 12, 2012

**Figure 10. City of Austin Pneumonia and Influenza Mortality:
2010-2013**



Data source: Center for Disease Control and Prevention 122 Cities Mortality: <http://wonder.cdc.gov/mmwr/mmwrmort.asp>


Austin/Travis County influenza surveillance does not attempt to capture all cases of influenza or influenza-like illness. These data should be used for trending purposes over time and for identifying types/strains of influenza that are occurring in the Austin area rather than for estimating the total number of cases.



**Austin/Travis County Health and Human Services Department
Epidemiology and Health Statistics Unit**




**Figure 11. RODS Syndromic Surveillance for the Respiratory Prodrome Category,
Travis County: 2010-2013**



Data source: University of Pittsburgh Real-time Outbreak and Disease Surveillance System. Data is an aggregation of Austin area chief complaint hospital data

**Figure 12. RODS Syndromic Surveillance for Influenza-Like-Illness,
Travis County: 2010-2013**



Data source: University of Pittsburgh Real-time Outbreak and Disease Surveillance System. Data is an aggregation of Austin area chief complaint hospital data

Austin/Travis County influenza surveillance does not attempt to capture all cases of influenza or influenza-like illness. These data should be used for trending purposes over time and for identifying types/strains of influenza that are occurring in the Austin area rather than for estimating the total number of cases.



Austin/Travis County Health and Human Services Department Epidemiology and Health Statistics Unit



- 1 MMWR is the Morbidity and Mortality weekly report week published by the CDC.
- 2 PCR testing is performed for specimens referred by area sentinel Influenza surveillance reporters. Providers interested in becoming sentinel providers may call the Austin/Travis County Health and Human Services Epidemiology and Health Statistics Unit at (512) 972-5555.
- 3 Influenza A is commonly split into 2 subtypes: H1N1 and H3N2. Both strains can circulate each Influenza season.
- 4 Influenza is not a reportable condition in Texas; therefore, data is provided by sentinel surveillance reporters and is only a sample of the Influenza activity occurring in the Austin/Travis County area.
- 5 Data represent rapid Influenza testing; these tests provide quick results reporting only Influenza A or B (no subtyping). Early in the flu season, results should be used with caution due to false positive results which can occur, especially during times when Influenza activity is low.
- 6 Influenza data is collected from a variety of reporters; the number of reporters can vary from week to week.
- 7 Non-Differentiated refers to rapid test results that do not differentiate between Influenza A and B.
- 8 Data for Austin/Travis County ILI reporters only; the number of reporters can vary from week to week.
- 9 Influenza is confirmed via PCR testing and a further subtyping may be performed. PCR testing is performed for specimens referred by area Influenza surveillance reporters.
- 10 Positive laboratory results are reported according to specimen collection date or date received in the lab if the former is unknown.
- 11 Antigenic characterization and antiviral resistance is obtained from the Texas Influenza Surveillance Report <http://www.dshs.state.tx.us/idcu/disease/Influenza/surveillance/2013/>.
- 12 Influenza activity level corresponds to the current MMWR week only and does not reflect the previous weeks' activity.
- 13 The majority of Influenza cases are not reportable by law to the Texas Department of State Health Services. This map contains data from sentinel sites and does not represent all Influenza cases in the state.
- 14 Laboratory data in 2013-2014 season reports may not be comparable to reports from previous seasons because of the inclusion of DSHS and LRN laboratory data for the current season.
- 15 Some non-NREVSS reporters also contribute to the RSV data.
- 16 Percentages based on the total number of specimens B positive and subtyped A specimens.